arXiv:1609.07481v1 [math.CA] 23 Sep 2016

Diefferential equations satisfied by $a(q) = \sum_{m,n\in\mathbb{Z}} q^{m^2+mn+n^2}$

By Kazuhide Matsuda

Faculty of Fundamental Science, National Institute of Technology, Niihama College,
7-1 Yagumo-chou, Niihama, Ehime, Japan, 792-8580.
E-mail: matsuda@sci.niihama-nct.ac.jp
Fax: 81-0897-37-7809

Abstract In this paper, we deive a differential equation satisfied by the following modular form of level three,

$$a(q) = \sum_{m,n\in\mathbb{Z}} q^{m^2+mn+n^2},\ q = \exp(2\pi i\tau),\ \tau \in \mathbb{H}^2,$$

where $\mathbb{H}^2$ is the upper half plane.
Key Words: theta function; theta constant; rational characteristics; Hecke group.
MSC(2010) 14K25; 11E25

1 Introduction

Throughout this paper, let $\mathbb{N}_0$, $\mathbb{N}$, denote the set of nonnegative integers, positive integers. For the positive integers j, k, and $n \in \mathbb{N}$, $d_{j,k}(n)$ denotes the number of positive divisors d of n such that $d \equiv j \bmod k$. Moreover, For k and $n \in \mathbb{N}$, $\sigma_k(n)$ is the sum of k-th power of positive divisors of n, and $d_{j,k}(n) = \sigma_k(n) = 0$ for $n \in \mathbb{Q} \setminus \mathbb{N}_0$. For each $n \in \mathbb{N}$, set

$$\left(\frac{n}{3}\right) = \begin{cases} +1, & \text{if } n \equiv 1 \bmod 3, \\ -1, & \text{if } d \equiv -1 \bmod 3, \\ 0, & \text{if } d \equiv 0 \bmod 3. \end{cases}$$

The *upper half plane* $\mathbb{H}^2$ is defined by $\mathbb{H}^2 = \{\tau \in \mathbb{C} \mid \Im\tau > 0\}$. Throughout this paper, set $q = \exp(2\pi i\tau)$ and defione the *Dedekind eta function* by

$$\eta(\tau) = q^{\frac{1}{24}} \prod_{n=1}^{\infty}(1-q^n) = q^{\frac{1}{24}}(q;q)_\infty.$$

The Eisenstein series E_2, E_4 and E_6 are defined by

$$E_2(q) = E_2(\tau) := 1 - 24\sum_{n=1}^{\infty}\sigma(n)q^n,$$
$$E_4(q) = E_4(\tau) := 1 + 240\sum_{n=1}^{\infty}\sigma_3(n)q^n,$$
$$E_6(q) = E_4(\tau) := 1 - 504\sum_{n=1}^{\infty}\sigma_5(n)q^n.$$

The aim of this paper is to derive differential equations satisfied by

$$a(q) = a(\tau) := \sum_{m,n\in\mathbb{Z}} q^{m^2+mn+n^2} = 1 + 6\sum_{n=1}^{\infty}(d_{1,3}(n) - d_{2,3}(n))q^n,\ q = \exp(2\pi i\tau).$$

For the proof of the second equality, see Berndt [3, pp. 79] or Dickson [5, pp. 68].

There were many researches in ordinary differential equations (ODEs) satisfied by modular forms. Classical examples include Ramanujan's coupled ODEs for Eisenstein series E_2, E_4, E_6; Rankin's fourth order ODE satisfied by $\Delta = \eta^{24}(\tau)$; Jacobi's third order ODE satisfied by the theta null functions $\vartheta_2, \vartheta_3, \vartheta_4$, which are defined in equation (2.1). For the proof, see Berndt [3, pp.92], Rankin [18] and Jacobi [11].

Halphen [10] rewrote Jacobi's ODE as a nonliear dynamical system:

$$X' + Y' = 2XY,\ Y' + Z' = 2YZ,\ Z' + X' = 2ZX.$$

ODEs of quadratic type are called Halphen type systems.

Recently, Ohyama [15, 16] reconsidered Jacobi's ODE from Picard-Fuchs equations of elliptic modular surfaces, and following Jacobi's idea, derived a Halphen type system satisfied by modular forms of level three. Following Ohyama, Mano [14] derived ODEs satisfied by modular forms of level five.

Ramanujan's ODEs are

$$q\frac{E_2}{dq} = \frac{(E_2)^2 - E_4}{12},\ q\frac{E_4}{dq} = \frac{E_2E_4 - E_6}{3},\ q\frac{E_6}{dq} = \frac{E_2E_6 - (E_4)^2}{2},$$

which is equivalent to Chazy's third order nonlinear ODE,

$$y''' = 2yy'' - 3(y')^2.$$

Ramamani [17] introduced

$$\mathcal{P} = 1 - 8\sum_{n=1}^{\infty}\frac{(-1)^n nq^n}{1-q^n},\ \tilde{\mathcal{P}} = 1 + 24\sum_{n=1}^{\infty}\frac{nq^n}{1+q^n},\ \mathcal{Q} = 1 + 16\sum_{n=1}^{\infty}\frac{(-1)^n n^3q^n}{1-q^n},$$

and derived ODEs satisfied by $\mathcal{P}, \tilde{\mathcal{P}}, \mathcal{Q}$, modular forms of $\Gamma_0(2)$. Ablowitz [1] et al. showed that these ODEs are equivalent to third order nonlinear ODE found by Bureau [4], and equivalent to a Halphen type system. Maiser [13] generalized these results to the Hecke group $\Gamma_0(N)$ $(N = 2, 3, 4)$.

Let us define the operators

$$\theta := q\frac{d}{dq} = \frac{1}{2\pi i}\frac{d}{d\tau},\ \partial := \partial_k = 12\theta - kE_2(q),\ (k = 1, 2, 3, \ldots).$$

For the properties of the operators ∂, see Kilford [6, pp. 35-37], or Rankin [18].

Our main theorem is as follows.

Theorem 1.1. *For $q \in \mathbb{C}$ with $|q| < 1$, define the operator by*

$$\partial = 12q\frac{d}{dq} - E_2(q).$$

Then, we have

$$\partial a(q) = 3a^3(q) - 4\frac{(q;q)^9}{(q^3;q^3)^3}, \tag{1.1}$$

and

$$\left(4q\frac{d}{dq} - E_2(q^3)\right) a(q) = -a^3(q) + 36q\frac{(q^3;q^3)^9}{(q;q)^3}, \tag{1.2}$$

where

$$\frac{(q;q)^9}{(q^3;q^3)^3} = 1 - 9\sum_{n=1}^{\infty} q^n \left(\sum_{d|n} d^2 \left(\frac{d}{3}\right)\right), \tag{1.3}$$

and

$$q\frac{(q^3;q^3)^9}{(q;q)^3} = \sum_{n=1}^{\infty} q^n \left(\sum_{d|n} d^2 \left(\frac{n/d}{3}\right)\right). \tag{1.4}$$

Section 2 reviews Farkas and Krs's theory of theta functions with rational characteristics. Section 3 treats some theta functional formulas. In particular, we prove equations (1.3) and (1.4). Section 4 proves Theorem 1.1. Section 5 shows some Ramanujan's identity of $a(q)$, which expresses $a(q)$ by the Dedekind's eta functions. Section 6 derives more product-series identities.

2 The properties of the theta functions

2.1 Definitions

Following Farkas and Kra [8], we introduce the *theta function with characteristics,* which is defined by

$$\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](\zeta,\tau)=\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](\zeta):=\sum_{n\in\mathbb{Z}}\exp\left(2\pi i\left[\frac{1}{2}\left(n+\frac{\epsilon}{2}\right)^2\tau+\left(n+\frac{\epsilon}{2}\right)\left(\zeta+\frac{\epsilon'}{2}\right)\right]\right),$$

where $\epsilon,\epsilon'\in\mathbb{R}$, $\zeta\in\mathbb{C}$, and $\tau\in\mathbb{H}^2$. The *theta constants* are given by

$$\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right]:=\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](0,\tau).$$

In particular, note that

$$\vartheta_2=\theta\left[\begin{array}{c}1\\ 0\end{array}\right],\ \vartheta_3=\theta\left[\begin{array}{c}0\\ 0\end{array}\right],\ \vartheta_4=\theta\left[\begin{array}{c}0\\ 1\end{array}\right]. \tag{2.1}$$

Furthermore, we denote the derivative coefficients of the theta function by

$$\theta'\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right]:=\frac{\partial}{\partial\zeta}\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](\zeta,\tau)\Big|_{\zeta=0},\ \theta''\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right]:=\frac{\partial^2}{\partial\zeta^2}\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](\zeta,\tau)\Big|_{\zeta=0}.$$

In particular, Jacobi's derivative formula is given by

$$\theta'\left[\begin{array}{c}1\\ 1\end{array}\right]=-\pi\theta\left[\begin{array}{c}0\\ 0\end{array}\right]\theta\left[\begin{array}{c}1\\ 0\end{array}\right]\theta\left[\begin{array}{c}0\\ 1\end{array}\right]. \tag{2.2}$$

2.2 Basic properties

We first note that for $m,n\in\mathbb{Z}$,

$$\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](\zeta+n+m\tau,\tau)=\exp(2\pi i)\left[\frac{n\epsilon-m\epsilon'}{2}-m\zeta-\frac{m^2\tau}{2}\right]\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](\zeta,\tau), \tag{2.3}$$

and

$$\theta\left[\begin{array}{c}\epsilon+2m\\ \epsilon'+2n\end{array}\right](\zeta,\tau)=\exp(\pi i\epsilon n)\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](\zeta,\tau). \tag{2.4}$$

Furthermore, it is easy to see that

$$\theta\left[\begin{array}{c}-\epsilon\\ -\epsilon'\end{array}\right](\zeta,\tau)=\theta\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](-\zeta,\tau)\text{ and }\theta'\left[\begin{array}{c}-\epsilon\\ -\epsilon'\end{array}\right](\zeta,\tau)=-\theta'\left[\begin{array}{c}\epsilon\\ \epsilon'\end{array}\right](-\zeta,\tau).$$

For $m, n \in \mathbb{R}$, we see that

$$\theta \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right] \left(\zeta + \frac{n+m\tau}{2}, \tau \right) = \exp(2\pi i) \left[-\frac{m\zeta}{2} - \frac{m^2\tau}{8} - \frac{m(\epsilon'+n)}{4} \right] \theta \left[\begin{array}{c} \epsilon+m \\ \epsilon'+n \end{array} \right] (\zeta, \tau). \tag{2.5}$$

We note that $\theta \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right] (\zeta, \tau)$ has only one zero in the fundamental parallelogram, which is given by

$$\zeta = \frac{1-\epsilon}{2}\tau + \frac{1-\epsilon'}{2}.$$

2.3 Jacobi's triple product identity

All the theta functions have infinite product expansions, which are given by

$$\theta \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right] (\zeta, \tau) = \exp\left(\frac{\pi i \epsilon \epsilon'}{2} \right) x^{\frac{\epsilon^2}{4}} z^{\frac{\epsilon}{2}} \times \prod_{n=1}^{\infty} (1-x^{2n})(1+e^{\pi i \epsilon'} x^{2n-1+\epsilon} z)(1+e^{-\pi i \epsilon'} x^{2n-1-\epsilon}/z), \tag{2.6}$$

where $x = \exp(\pi i \tau)$ and $z = \exp(2\pi i \zeta)$. Therefore, it follows from Jacobi's derivative formula (2.2) that

$$\theta' \left[\begin{array}{c} 1 \\ 1 \end{array} \right] (0, \tau) = -2\pi q^{\frac{1}{8}} \prod_{n=1}^{\infty} (1-q^n)^3, \ q = \exp(2\pi i \tau).$$

2.4 Spaces of N-th order θ-functions

Following Farkas and Kra [8], we define $\mathcal{F}_N \left[\begin{array}{c} \epsilon \\ \epsilon' \end{array} \right]$ to be the set of entire functions f that satisfy the two functional equations,

$$f(\zeta+1) = \exp(\pi i \epsilon)\ f(\zeta),$$

and

$$f(\zeta+\tau) = \exp(-\pi i)[\epsilon' + 2N\zeta + N\tau]\ f(\zeta), \quad \zeta \in \mathbb{C},\ \tau \in \mathbb{H}^2,$$

where N is a positive integer and $\begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} \in \mathbb{R}^2$. This set of functions is called the space of *N-th order θ-functions with characteristic* $\begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix}$. Note that

$$\dim \mathcal{F}_N \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} = N.$$

For its proof, see Farkas and Kra [8, pp.133].

2.5 The heat equation

The theta function satisfies the following heat equation:

$$\frac{\partial^2}{\partial \zeta^2} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau) = 4\pi i \frac{\partial}{\partial \tau} \theta \begin{bmatrix} \epsilon \\ \epsilon' \end{bmatrix} (\zeta, \tau). \tag{2.7}$$

3 Some theta functional formulas

Theorem 3.1. *For every $x, y \in \mathbb{C}$, the following holds:*

$$\frac{d^2}{dx^2} \log \theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} (x) - \frac{d^2}{dy^2} \log \theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} (y) = \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^2 \frac{\theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} (x-y) \theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} (x+y)}{\theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (x) \theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (y)}. \tag{3.1}$$

Proof. For the proof, see Liu [12]. □

Corollary 3.2. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{\eta^{10}(3\tau)}{\eta^3(9\tau)\eta^3(\tau)} = 1 + 3\sum_{n=1}^{\infty} q^n \left(\sum_{\substack{d \mid n \\ (\frac{n}{d}, 3) = 1}} d \right) + 3\sum_{n=1}^{\infty} q^{3n} \left(\sum_{\substack{d \mid n \\ (d, 3) = 1}} d \right) - 18\sum_{n=1}^{\infty} \sigma(n) q^{9n}, \tag{3.2}$$

and

$$\frac{\eta^3(\tau)\eta^3(9\tau)}{\eta^2(3\tau)} = \sum_{n=1}^{\infty} q^n \left(\sum_{\substack{d \mid n \\ \frac{n}{d} \equiv 1 \mod 3}} d\left(\frac{d}{3}\right) \right) - \sum_{n=1}^{\infty} q^n \left(\sum_{\substack{d \mid n \\ \frac{n}{d} \equiv 2 \mod 3}} d\left(\frac{d}{3}\right) \right), \tag{3.3}$$

where $q = \exp(2\pi i \tau)$.

Proof. The corollary can be proved by substituting $(x,y) = (-\tau/3, -1/3)$ or $(x,y) = (-\tau/3 - 1/3, -\tau/3 + 1/3)$ and applying Jacobi's derivative formula (2.6). □

Theorem 3.3. *For every $z \in \mathbb{C}$, we have*

$$\frac{d^3}{dz^3} \log \theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z) = \theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}^3 \frac{\theta \begin{bmatrix} 1 \\ 1 \end{bmatrix} (2z)}{\theta^4 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z)}. \tag{3.4}$$

Proof. The theorem can be proved by dividing both sides of equation (3.1) by $(x-y)$ and taking the limit $x \longrightarrow y$. □

Corollary 3.4. *For every $\tau \in \mathbb{H}^2$, we have*

$$\frac{\eta^9(\tau)}{\eta^3(3\tau)} = 1 - 9 \sum_{n=1}^{\infty} q^n \left(\sum_{d|n} d^2 \left(\frac{d}{3} \right) \right) \tag{3.5}$$

and

$$\frac{\eta^9(3\tau)}{\eta^3(\tau)} = \sum_{n=1}^{\infty} q^n \left(\sum_{d|n} d^2 \left(\frac{n/d}{3} \right) \right), \tag{3.6}$$

where $q = \exp(2\pi i \tau)$.

Proof. The corollary can be proved by substituting $z = -1/3$ or $z = -\tau/3$ in equation (3.4) and applying Jacobi's triple product identity (2.6). □

4 Proof of Theorem 1.1

4.1 Preliminary results

Proposition 4.1. *For every $\tau \in \mathbb{H}^2$, we have*

$$3 \frac{\theta'' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 6 \left\{ \frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} \right\}^2 = 0, \tag{4.1}$$

$$3 \frac{\theta'' \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix}} - \frac{\theta''' \begin{bmatrix} 1 \\ 1 \end{bmatrix}}{\theta' \begin{bmatrix} 1 \\ 1 \end{bmatrix}} + 6 \left\{ \frac{\theta' \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix}} \right\}^2 = 0. \tag{4.2}$$

$$3\frac{\theta''\begin{bmatrix}\frac{1}{3}\\ \frac{1}{3}\end{bmatrix}}{\theta\begin{bmatrix}\frac{1}{3}\\ \frac{1}{3}\end{bmatrix}}-\frac{\theta'''\begin{bmatrix}1\\ 1\end{bmatrix}}{\theta'\begin{bmatrix}1\\ 1\end{bmatrix}}+6\left\{\frac{\theta'\begin{bmatrix}\frac{1}{3}\\ \frac{1}{3}\end{bmatrix}}{\theta\begin{bmatrix}\frac{1}{3}\\ \frac{1}{3}\end{bmatrix}}\right\}^2=0, \tag{4.3}$$

and

$$3\frac{\theta''\begin{bmatrix}\frac{1}{3}\\ \frac{5}{3}\end{bmatrix}}{\theta\begin{bmatrix}\frac{1}{3}\\ \frac{5}{3}\end{bmatrix}}-\frac{\theta'''\begin{bmatrix}1\\ 1\end{bmatrix}}{\theta'\begin{bmatrix}1\\ 1\end{bmatrix}}+6\left\{\frac{\theta'\begin{bmatrix}\frac{1}{3}\\ \frac{5}{3}\end{bmatrix}}{\theta\begin{bmatrix}\frac{1}{3}\\ \frac{5}{3}\end{bmatrix}}\right\}^2=0. \tag{4.4}$$

Proof. Consider the following elliptic function:

$$\varphi_1(z)=\frac{\theta^3\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}(z)}{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)},\quad \varphi_2(z)=\frac{\theta^3\begin{bmatrix}\frac{1}{3}\\ 1\end{bmatrix}(z)}{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)},\quad \varphi_3(z)=\frac{\theta^3\begin{bmatrix}\frac{1}{3}\\ \frac{1}{3}\end{bmatrix}(z)}{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)},\quad \varphi_4(z)=\frac{\theta^3\begin{bmatrix}\frac{1}{3}\\ \frac{5}{3}\end{bmatrix}(z)}{\theta^3\begin{bmatrix}1\\ 1\end{bmatrix}(z)}.$$

In the fundamental parallelogram, the pole of $\varphi_j(z)$ $(j=1,2,3,4)$ is $z=0$, which implies that $\mathrm{Res}(\varphi_j(z),0)=0$. Therefore, the proposition follows. □

Proposition 4.2. *For every $\tau\in\mathbb{H}^2$, we have*

$$\frac{\theta'''\begin{bmatrix}1\\ 1\end{bmatrix}}{\theta'\begin{bmatrix}1\\ 1\end{bmatrix}}=4\pi i\frac{d}{d\tau}\log\theta'\begin{bmatrix}1\\ 1\end{bmatrix}=-\pi^2E_2(q),\ q=\exp(2\pi i\tau).$$

Proof. The proposition follows from Jacobi's triple product identity (2.6). □

Proposition 4.3. For every $\tau\in\mathbb{H}^2$, we have

$$\frac{\theta'\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}}{\theta\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}}=-\frac{\pi}{\sqrt{3}}a(\tau),\ \textit{and}\ \frac{\theta'\begin{bmatrix}\frac{1}{3}\\ 1\end{bmatrix}}{\theta\begin{bmatrix}\frac{1}{3}\\ 1\end{bmatrix}}=\frac{\pi i}{3}a(\tau/3). \tag{4.5}$$

Proof. The proposition follows from Jacobi's triple product identity (2.6). □

4.2 Proof of equation (1.1)

Proof. By substituting $z = -1/3$ in equation (3.4), we have

$$\frac{\theta'\begin{bmatrix}1\\1\end{bmatrix}^3}{\theta^3\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}} = 4\pi i\frac{d}{d\tau}\left\{\frac{\theta'\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}}{\theta\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}}\right\} - 2\frac{\theta''\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}}{\theta\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}}\frac{\theta'\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}}{\theta\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}} + 2\left\{\frac{\theta'\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}}{\theta\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}}\right\}^3 = 0. \tag{4.6}$$

The theorem follows from Propositions 4.1, 4.2 and 4.3. □

4.3 Proof of equation (1.2)

Proof. By substituting $z = -\tau/3$ in equation (3.4), we have

$$-\frac{\theta'\begin{bmatrix}1\\1\end{bmatrix}^3}{\theta^3\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}} = 4\pi i\frac{d}{d\tau}\left\{\frac{\theta'\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}}{\theta\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}}\right\} - 2\frac{\theta''\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}}{\theta\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}}\frac{\theta'\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}}{\theta\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}} + 2\left\{\frac{\theta'\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}}{\theta\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}}\right\}^3 = 0. \tag{4.7}$$

The theorem can be obtained by applying Propositions 4.1, 4.2 and 4.3 and changing $\tau \longrightarrow 3\tau$. □

5 Some Ramanujan's identity

5.1 Farkas and Kra's cubic identity

Theorem 5.1. (Farkas and Kra [8, pp. 193]) *For every $\tau \in \mathbb{H}^2$, we have*

$$\theta^3\begin{bmatrix}\frac{1}{3}\\\frac{1}{3}\end{bmatrix} + \theta^3\begin{bmatrix}\frac{1}{3}\\\frac{5}{3}\end{bmatrix} = \theta^3\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}, \tag{5.1}$$

and

$$\exp\left(\frac{\pi i}{3}\right)\theta^3\begin{bmatrix}\frac{1}{3}\\\frac{1}{3}\end{bmatrix} + \exp\left(\frac{2\pi i}{3}\right)\theta^3\begin{bmatrix}\frac{1}{3}\\\frac{5}{3}\end{bmatrix} = \theta^3\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}. \tag{5.2}$$

Proof. Consider the following elliptic functions:

$$\varphi(z) = \frac{\theta^3\begin{bmatrix}1\\1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{1}{3}\\\frac{1}{3}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}(z)\theta\begin{bmatrix}\frac{1}{3}\\\frac{5}{3}\end{bmatrix}(z)}, \quad \psi(z) = \frac{\theta^3\begin{bmatrix}1\\1\end{bmatrix}(z)}{\theta\begin{bmatrix}\frac{1}{3}\\\frac{1}{3}\end{bmatrix}(z)\theta\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}(z)\theta\begin{bmatrix}\frac{5}{3}\\\frac{1}{3}\end{bmatrix}(z)}.$$

By $\varphi(z)$, we prove equation (5.1). Equation (5.2) can be obtained by $\psi(z)$ in the same way.

Note that in the fundamental parallelogram, the poles of $\varphi(z)$ are $(\tau+1)/3$, $\tau/3$, and $(\tau-1)/3$. Direct computations yield

$$\mathrm{Res}\left(\varphi(z), \frac{\tau+1}{3}\right) = -\frac{\theta^3\begin{bmatrix}\frac{1}{3}\\ \frac{1}{3}\end{bmatrix}}{\theta'\begin{bmatrix}1\\1\end{bmatrix}\theta^2\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}}, \quad \mathrm{Res}\left(\varphi(z), \frac{\tau}{3}\right) = \frac{\theta^3\begin{bmatrix}\frac{1}{3}\\ 1\end{bmatrix}}{\theta'\begin{bmatrix}1\\1\end{bmatrix}\theta^2\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}},$$

and

$$\mathrm{Res}\left(\varphi(z), \frac{\tau-1}{3}\right) = -\frac{\theta^3\begin{bmatrix}\frac{1}{3}\\ \frac{5}{3}\end{bmatrix}}{\theta'\begin{bmatrix}1\\1\end{bmatrix}\theta^2\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}}.$$

From the residue theorem, it follows that

$$\mathrm{Res}\left(\varphi(z), \frac{\tau+1}{3}\right) + \mathrm{Res}\left(\varphi(z), \frac{\tau}{3}\right) + \mathrm{Res}\left(\varphi(z), \frac{\tau-1}{3}\right) = 0,$$

which implies equation (5.1). □

5.2 Ramanujan's identity

Theorem 5.2. (Ramanujan [2, pp. 346]) *For every $\tau \in \mathbb{H}^2$, we have*

$$a(q) = \frac{\eta^3(\tau/3) + 3\eta(3\tau)}{\eta(\tau)}.$$

Proof. From Farkas [7], we recall the following identity:

$$\begin{aligned}&\frac{6\theta'\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}(0,\tau)}{\zeta_6\theta^3\begin{bmatrix}\frac{1}{3}\\ \frac{1}{3}\end{bmatrix}(0,\tau) + \theta^3\begin{bmatrix}\frac{1}{3}\\ 1\end{bmatrix}(0,\tau) + \zeta_6^5\theta^3\begin{bmatrix}\frac{1}{3}\\ \frac{5}{3}\end{bmatrix}(0,\tau)}\\ =&\frac{2\pi i q^{\frac{1}{12}}}{\prod_{n=0}^{\infty}(1-q^{3n+1})(1-q^{3n+2})} = 2\pi i\frac{e^{\frac{\pi i}{6}}}{\sqrt{3}}\frac{\theta\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}(0,\tau)}{\theta\begin{bmatrix}\frac{1}{3}\\ 1\end{bmatrix}(0,3\tau)},\end{aligned} \tag{5.3}$$

where $q = \exp(2\pi i\tau)$ and $\zeta_6 = \exp(2\pi i/6)$.

Theorem 5.1 yields that

$$\theta^3 \begin{bmatrix} \frac{1}{3} \\ \frac{1}{3} \end{bmatrix} = -\exp\left(\frac{2\pi i}{3}\right) \theta^3 \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} + \theta^3 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix},$$

and

$$\theta^3 \begin{bmatrix} \frac{1}{3} \\ \frac{5}{3} \end{bmatrix} = \exp\left(\frac{\pi i}{3}\right) \theta^3 \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} - \theta^3 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix},$$

which imply

$$\frac{\theta' \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}} = \frac{\pi \exp(\frac{2\pi i}{3})}{3\sqrt{3}} \times \frac{3\theta^3 \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} + \sqrt{3}i\theta^3 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix}}{\theta \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} (0, 3\tau)}.$$

Therefore, the theorem follows from Proposition 4.3 and Jacobi's triple product identity (2.6). □

6 More product-series identities

6.1 Some theta functional formulas

Proposition 6.1. *For every $(z, \tau) \in \mathbb{C} \times \mathbb{H}^2$, we have*

$$\theta^2 \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} \theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (z)\theta \begin{bmatrix} 1 \\ \frac{5}{3} \end{bmatrix} (z) + \exp\left(\frac{\pi i}{3}\right) \theta^2 \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} \theta \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} (z)\theta \begin{bmatrix} \frac{5}{3} \\ 1 \end{bmatrix} (z)$$
$$- \theta \begin{bmatrix} \frac{1}{3} \\ \frac{1}{3} \end{bmatrix} \theta \begin{bmatrix} \frac{1}{3} \\ \frac{5}{3} \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z) = 0, \tag{6.1}$$

$$\theta^2 \begin{bmatrix} \frac{1}{3} \\ \frac{5}{3} \end{bmatrix} \theta \begin{bmatrix} \frac{1}{3} \\ \frac{1}{3} \end{bmatrix} (z)\theta \begin{bmatrix} \frac{5}{3} \\ \frac{5}{3} \end{bmatrix} (z) + \theta^2 \begin{bmatrix} \frac{1}{3} \\ \frac{1}{3} \end{bmatrix} \theta \begin{bmatrix} \frac{1}{3} \\ \frac{5}{3} \end{bmatrix} (z)\theta \begin{bmatrix} \frac{5}{3} \\ \frac{1}{3} \end{bmatrix} (z)$$
$$- \exp\left(\frac{2\pi i}{3}\right) \theta \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} \theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} \theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z) = 0. \tag{6.2}$$

Proof. We prove equation (6.1). Equation (6.2) can be proved in the same way. We first note that $\dim \mathcal{F}_2 \begin{bmatrix} 0 \\ 0 \end{bmatrix} = 2$, and

$$\theta \begin{bmatrix} 1 \\ \frac{1}{3} \end{bmatrix} (z, \tau)\theta \begin{bmatrix} 1 \\ \frac{5}{3} \end{bmatrix} (z, \tau),\ \theta \begin{bmatrix} \frac{1}{3} \\ 1 \end{bmatrix} (z, \tau)\theta \begin{bmatrix} \frac{5}{3} \\ 1 \end{bmatrix} (z, \tau),\ \theta^2 \begin{bmatrix} 1 \\ 1 \end{bmatrix} (z, \tau) \in \mathcal{F}_2 \begin{bmatrix} 0 \\ 0 \end{bmatrix}.$$

Therefore, there exist some complex numbers, x_1, x_2 and x_3 not all zero such that

$$x_1\theta\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}(z,\tau)\theta\begin{bmatrix}1\\ \frac{5}{3}\end{bmatrix}(z,\tau)+x_2\theta\begin{bmatrix}\frac{1}{3}\\ 1\end{bmatrix}(z,\tau)\theta\begin{bmatrix}\frac{5}{3}\\ 1\end{bmatrix}(z,\tau)+x_3\theta^2\begin{bmatrix}1\\ 1\end{bmatrix}(z,\tau)=0.$$

Note that in the fundamental parallelogram, the zero of $\theta\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}(z)$, $\theta\begin{bmatrix}\frac{1}{3}\\ 1\end{bmatrix}(z)$, or $\theta\begin{bmatrix}1\\ 1\end{bmatrix}(z)$ is $z=1/3$, $\tau/3$ or 0. Substituting $z=1/3, \tau/3$, and 0, we have

$$\begin{aligned}
&&x_2\exp\left(-\frac{\pi i}{3}\right)\theta\begin{bmatrix}\frac{1}{3}\\ \frac{1}{3}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{3}\\ \frac{5}{3}\end{bmatrix}+x_3\theta^2\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}&=0,\\
x_1\theta\begin{bmatrix}\frac{1}{3}\\ \frac{1}{3}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{3}\\ \frac{5}{3}\end{bmatrix}&&+x_3\theta^2\begin{bmatrix}\frac{1}{3}\\ 1\end{bmatrix}&=0,\\
-x_1\theta^2\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix}\quad&+x_2\exp\left(-\frac{\pi i}{3}\right)\theta^2\begin{bmatrix}\frac{1}{3}\\ 1\end{bmatrix}&&=0.
\end{aligned}$$

Solving this system of equations, we have

$$(x_1,x_2,x_3)=\alpha\left(\theta^2\begin{bmatrix}\frac{1}{3}\\ 1\end{bmatrix},\exp\left(\frac{\pi i}{3}\right)\theta^2\begin{bmatrix}1\\ \frac{1}{3}\end{bmatrix},-\theta\begin{bmatrix}\frac{1}{3}\\ \frac{1}{3}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{3}\\ \frac{5}{3}\end{bmatrix}\right)\ \text{ for some } \alpha\in\mathbb{C}\setminus\{0\},$$

which proves the proposition. □

6.2 Product-series identities

Theorem 6.2. *For every $\tau\in\mathbb{H}^2$, we have*

$$\frac{\eta^{10}(3\tau)}{\eta^3(\tau)\eta^3(9\tau)}=1+3\sum_{n=1}^{\infty}(\sigma(n)-9\sigma(n/9))q^n, \tag{6.3}$$

and

$$\frac{\eta^3(\tau)\eta^3(9\tau)}{\eta^2(3\tau)}=\sum_{n=1}^{\infty}q^n\left(\sum_{\substack{d\mid n\\ \frac{n}{d}\equiv 1 \mod 3}}\frac{n}{d}\left(\frac{d}{3}\right)\right)-\sum_{n=1}^{\infty}q^n\left(\sum_{\substack{d\mid n\\ \frac{n}{d}\equiv 2 \mod 3}}\frac{n}{d}\left(\frac{d}{3}\right)\right), \tag{6.4}$$

where $q=\exp(2\pi i\tau)$.

Proof. By equations (4.1), (4.2), and (6.1), we derive equation (6.3). Equation (6.4) can be proved in the same way.

Comparing the coefficients of the term z^2 in equation (6.1), we have

$$\left\{\theta'\begin{bmatrix}1\\1\end{bmatrix}\right\}^2\frac{\theta\begin{bmatrix}\frac{1}{3}\\\frac{1}{3}\end{bmatrix}\theta\begin{bmatrix}\frac{1}{3}\\\frac{5}{3}\end{bmatrix}}{\theta^2\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}\theta^2\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}}=\frac{3}{2}\left(\frac{\theta''\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}}{\theta\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}}-\frac{\theta''\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}}{\theta\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}}\right)$$
$$=6\pi i\frac{d}{d\tau}\log\frac{\theta\begin{bmatrix}\frac{1}{3}\\1\end{bmatrix}}{\theta\begin{bmatrix}1\\\frac{1}{3}\end{bmatrix}}.$$

Therefore, equation (6.3) can be obtained by Jacobi's triple product identity (2.6). □

References

[1] M. J. Ablowitz, S. Chakravarty, and H. Hahn, *Integrable systems and modular forms of level 2,* J. Phys. A **39** (2006), 15341-15353.

[2] B. Berndt, *Ramanujan's notebooks. Part III*, Springer-Verlag, New York, 1991.

[3] B. C. Berndt, *Number theory in the spirit of Ramanujan,* Stud. Math. Libr., **34** American Mathematical Society, Providence, RI, 2006.

[4] F. J. Bureau, *Sur des systémes différentiels non linéaires du troisiéme ordre et les équations différentielles non linéaires associées,* Acad. Roy. Belg. Bull. Cl. Sci. (5) **73** , (1987), 335353.

[5] L. E. Dickson, *Modern Elementary Theory of Numbers,* University of Chicago Press, Chicago, 1939.

[6] L. J. P. Kilford, *Modular forms. A classical and computational introduction,* Imperial College Press, London, 2008.

[7] H. M. Farkas, *Theta functions in complex analysis and number theory.* Surveys in number theory, Dev. Math., **17**, (2008), 57-87.

[8] H. M. Farkas and Irwin Kra, *Theta constants, Riemann surfaces and the modular group,* AMS Grad. Studies in Math. **37** (2001).

[9] J. Harnad, and J. McKay, *Modular solutions to equations of generalized Halphen type,* R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **456** (2000), 261-294.

[10] G. Halphen, *Sur une system déquations differ entielles,* C. R. Acad. Sci., Paris **92** (1881), 1101-1103.

[11] C.G.J. Jacobi, *Uber die Differentialgleichung welcher die Reihen* $1 \pm 2q + 2q^4 \pm 2q^4 +$ etc., $2q^{1/4} + 2q^{9/4} + 2q^{25/4} +$ etc. *Genuge leisten,* J. Reine Angew. Math. **36**, (1848), 97-112

[12] Zhi-Guo, Liu, *A three-term theta function identity and its applications,* Adv. Math. **195** (2005), 1-23.

[13] R. S. Maier, *Nonlinear differential equations satisfied by certain classical modular forms,* Manuscripta Math. **134** (2011), 1-42.

[14] T. Mano, Toshiyuki, *Differential relations for modular forms of level five*, J. Math. Kyoto Univ. **42**, (2002), 41-55.

[15] Y. Ohyama, *Systems of nonlinear differential equations related to second order linear equations,* Osaka J. Math. **33** (1996), 927-949.

[16] Y. Ohyama, *Differential equations for modular forms of level three,* Funkcial. Ekvac. **44** (2001), 377-389.

[17] V. Ramamani, *On some algebraic identities connected with Ramanujan's work,* Ramanujan International Symposium on Analysis, Macmillan of India, New Delhi (1989) 277-291

[18] R. A. Rankin, *The construction of automorphic forms from the derivatives of a given form,* J. Indian Math. Soc. **20** (1956), 103-116.